



PROCEEDINGS

Open Access

Gene polymorphism and protein of human pro- and anti-inflammatory cytokines in Chinese healthy subjects and chronic periodontitis patients

Wings TY Loo^{1,2}, Chang-bin Fan³, Lan-jun Bai⁴, Yuan Yue⁵, Yi-ding Dou⁶, Min Wang⁵, Hao Liang⁵, Mary NB Cheung⁶, Louis WC Chow^{1*}, Jin-le Li⁵, Ye Tian⁵, Liu Qing¹

From Organisation for Oncology and Translational Research (OOTR) 7th Annual Conference Hong Kong. 13-14 May 2011

Abstract

Background: Periodontal disease is thought to arise from the interaction of various factors, including the susceptibility of the host, the presence of pathogenic organisms, and the absence of beneficial species. The genetic factors may play a significant role in the risk of periodontal diseases. Cytokines initiate, mediate and control immune and inflammatory responses. The aim of this study is to compare genotypes and soluble protein of pro and anti-inflammatory cytokines (IL-1 α , IL-1 β , IL-6, IFN- γ , IL-10, TNF- α and IL-4) in subjects with or free of chronic periodontitis.

Methods: A total of 1,290 Chinese subjects were recruited to this clinical trial: 850 periodontally healthy controls and 440 periodontal patients. All subjects were free of systemic diseases. Oral examinations were performed, and the following parameters were recorded for each subject: supragingival/subgingival calculus, gingival recession, bleeding on probing (BOP), probing depth (PD), clinical attachment loss (CAL), gingival recession and tooth mobility. The peripheral blood samples were collected for genetic and enzyme linked immunosorbent assay (ELISA) analysis. Restriction enzymes were used for digestion of amplified fragments of IL-1 α , IL-1 β , IL-6, IFN- γ , IL-10, TNF- α and IL-4.

Results: The protein expressions of patient and control samples for IL-1 α , IL-1 β , IL-6, TNF- α , IFN- γ , IL-10, and IL-4 measured by ELISA confirmed a statistically significant difference ($p < 0.001$). The digestion of fragments of various genes showed that the pro-inflammatory cytokines IL-1 α and TNF- α , and the anti-inflammatory cytokines IL-4 and IL-10 demonstrated a correlation with chronic inflammation in patients (χ^2 : $p < 0.001$). The remaining genes investigated in patients and healthy subjects (IL-1 β , IL-6, IFN- γ and IL-10) did not show any significant difference.

Conclusions: The cytokine gene polymorphisms may be used as a marker for periodontitis susceptibility, clinical behaviour and severity. This detection offers early diagnosis and induction of prophylaxis to other family members against disease progression.

Background

Periodontal disease is thought to arise from the interaction of various factors, including the susceptibility of the host, the presence of pathogenic organisms and the absence of beneficial species [1]. Although bacteria

cause plaque-induced inflammatory periodontal diseases, progression and clinical characteristics of these diseases are influenced by both acquired and genetic factors that can modify susceptibility to infection [2] Reports have indicated that genetic factors may play a significant role in the risk of periodontal diseases [3,4].

Cytokines are soluble proteins which are secreted by cells to act as a messenger that transmits signals to other

* Correspondence: lwcchow@unimed.hk

¹UNIMED Medical Institute, Hong Kong SAR

Full list of author information is available at the end of the article

cells. They initiate, mediate and control immune and inflammatory responses; they also regulate growth and differentiation of cells [4]. Gingival epithelial cells produce a broad range of cytokines, among which, interleukin-1 α (IL-1 α), interleukin-1 β (IL-1 β), tumour necrosis factor- α (TNF- α), interleukin-6 (IL-6) and interferon- γ (IFN- γ) are the pro-inflammatory cytokines whereas interleukin-4 (IL-4) and interleukin-10 (IL-10) are the anti-inflammatory cytokines [5-7].

IL-1 consists of at least two separate gene products, IL-1 α and IL-1 β , which have common biological activities but limited homology at the nucleotide and peptide levels [8]. The gene encoding IL-1 is assigned to chromosome 2q13-21 [9]. The carriage of certain alleles of IL-1 α and IL-1 β is associated with the incidence and the severity of periodontal disease, in particular chronic periodontitis (CP), because these carriers produce more IL-1 in response to plaque than genotype negative individuals [2,10,11]. The presence of a single base-pair C-T transpositional polymorphism upstream of the IL-1 α gene at position -889 affects the production of the IL-1 α protein. The IL-1 α -889 T allele was associated with an almost four-fold increase in IL-1 α levels in gingival crevicular fluid [12]. A weak positive association was found concerning IL-1 β T(-511)C and chronic periodontal disease [13]. In other studies, no association was found for IL-1 α C(-889) T and -1 β T(-511)C with aggressive periodontitis [13,14].

TNF- α is located in 6p21.3 of chromosome 6 within the major histocompatibility complex [15]. Eight single nucleotide polymorphisms (SNPs) in the promoter region of this gene have been studied at positions -1031T/C, -863C/A, -857C/T, -575G/A, -376G/A, -308G/A, -244G/A, and -238G/A [16-19]. Many researchers investigated the possible link between the -308 polymorphism in the TNF- α gene and periodontitis because a G to A polymorphism at the -308 position of the TNF- α promoter region was suggested to influence TNF- α production and monocytes of patients with periodontitis [20,21]. Higher allele 2 frequency was observed in patients with chronic periodontitis than in healthy individuals [22]. Aggressive periodontitis was not linked to this locus [23-25]. The remaining promoter polymorphisms were not associated with periodontitis, apart from one study that found an association between the -1031, -863, -857 SNPs and chronic periodontitis severity in the Japanese population [23,26,27].

The IL-6 gene is assigned to chromosome 7p21. Various SNPs in the promoter region of this gene have been studied at positions -174G/C, -190C/T, -572C/G, -597G/A, -1363G/T, -1480C/G and -6106A/T [28,29]. In periodontitis, the G-C SNP at the -174 position correlated with chronic periodontitis susceptibility in Brazilians and Caucasians, but not the Japanese [30,31]. Periodontitis patients carrying one or two copies of the rare allele in the IL-6 -174 polymorphism displayed significantly higher serum

IL-6 and C-reactive protein concentrations [32]. Carriers of the rare allele at this position was associated with less reduction in probing depths among chronic periodontitis patients after delivery of standard non-surgical periodontal therapy [33]. On the contrary, another study found that the -174 R-allele carrier individuals have decreased plasma levels of IL-6 and lower IL-6 gene transcriptional activity compared with N/N individuals, suggesting that low IL-6 response may hamper an individual's defense against periodontal pathogens [34]. In a meta-analysis of cytokine gene polymorphisms of 53 studies, the IL-6 -174 polymorphism did not exhibit any association with chronic periodontitis (CP) [13].

The mRNA expression and/or concentration of IFN- γ in gingival crevicular fluid, gingival tissues, and serum were able to affect gingivitis, probing depths and alveolar bone loss [35]. Very few studies have investigated the connection of polymorphism of IFN- γ with periodontitis to date. Polymorphism in gene IFN- γ was found to be functionally relevant and causes differences in the immunoregulatory activity of its cytokine molecules. The T allele of the IFN- γ 874 T/A is found in high producers of IFN- γ [36]. No association was found in the only existing study dealing with the IFN- γ polymorphism 874T/A and CP [37].

The gene for IL-4 is localized in chromosome 5q31.1 [38]. The presence of IL-4-producing cells and the percentage of IL-4-expressing cells were significantly higher in established and severe periodontitis lesions than in gingivitis tissues. IL-4 levels in the serum of patients were higher in chronic periodontitis but these levels did not correlate with the degree of bone loss or pocket formation [39]. Mout *et al.* identified promoter SNP at position -590 and a 70-bp variable numbers of tandem repeat (VNTR) polymorphism at intron 2 [40]. However, many reports have failed to establish a connection between these loci and periodontal disease susceptibility and severity [41,42].

The gene encoding IL-10 was mapped to chromosome 1q31-32 [43]. The -1082 G/A locus was not associated with chronic periodontitis susceptibility in most Caucasian populations except in one Swedish study, but was linked to CP severity [44]. The -1082 single nucleotide polymorphism was associated with high *in vitro* IL-10 production [45]. There was a complete absence of the N-allele carriage at position -1082 among the Japanese in contrast to Caucasians where the -1082 N-allele is the most occurring variant [45,46].

The aim of this study is to compare genotypes and soluble protein (serum) of pro and anti-inflammatory cytokines in subjects with or free of chronic periodontitis.

Methods

Selection of subjects and clinical examination

From September 2004 to March 2007, a total of 1,290 subjects were recruited to this clinical trial: 850

periodontally healthy subjects and 440 periodontitis patients. Healthy subjects were selected from colleges and the community setting to be in the control group. They were invited for oral clinical examinations at West China Hospital of Stomatology, Sichuan University, PRC. Among the 850 healthy controls, 306 were female and 544 were male, with an average age of 42.9 years. 440 periodontal patients (180 females and 260 males, aged 28 to 65 years) with moderate to severe chronic periodontitis were recruited from West China Hospital of Stomatology, Sichuan University, PRC. The data of the recruited subjects are presented in Table 1.

The sample size of this study was determined based on the reports of Machin and Lemeshow. In the report, the sample size was calculated based on 0.05 level of significance for two arms to achieve 90% power [47,48]. The study protocols were approved by the Ethics Committee, West China Hospital of Stomatology, Sichuan University. Informed consent was obtained from all subjects.

The diagnosis of CP was made following the criteria defined by the American Academy of Periodontology in 1999 [49]. The periodontal disease classification system forms the basis for the criteria of diagnosis of periodontitis [2,49,50].

All subjects were free of oral soft tissue abnormalities or severe dental caries. An intra-oral examination of periodontal conditions including supragingival/subgingival calculus, gingival recession, bleeding on probing (BOP), probing depth (PD), clinical attachment loss (CAL), gingival recession and tooth mobility was performed. The diagnostic criteria for periodontitis were referred to the 1999 International Classification of the Periodontal Disease and Conditions [49]. Subjects who were examined and determined to be free of periodontal disease, gingival recession, CAL and probing depth of greater than 3 mm were regarded as periodontally healthy. If subjects presented with probing depth of

greater than 5 mm, CAL greater than 4 mm, some degree of gingival recession and tooth mobility, they were regarded to be severe chronic periodontitis patients [13,23,31].

Preparation of control and patient blood samples

Peripheral blood samples were collected by direct venipuncture from the arm vein of each subject: 20 ml in lithium heparin tubes and 10 ml in clot blood tubes (BD Vacutainer, NJ USA). The samples were centrifuged for 10 min at 1,500 rpm, and serum and plasma was then collected for enzyme linked immunosorbent assay (ELISA) analysis. The remaining cellular components were transferred to a 50 ml centrifuge tube and an additional 45 ml red blood cell lysis buffer was added. The tube was inverted several times and then centrifuged for 10 min at 1,500 rpm. The supernatants were discarded, and the remaining components were washed with 0.9% PBS used for DNA extraction.

Extraction of DNA from samples and polymerase chain reaction (PCR) to amplify the polymorphic site

Genomic DNA was extracted from blood samples following the blood protocols from the QIAamp DNA Blood Mini Kit (QIAGEN, MD, USA). The concentration of DNA was estimated by measurements of OD₂₆₀ by a spectrophotometer (U1800, Hitachi, Japan). Extracted DNA was labeled and stored at -80°C until use.

A PCR kit (Promega Corporation, U.S.A), consisting of nuclease-free water and PCR Master Mix was used according to the manufacturer's instructions. The PCR Master Mix includes 50 units/ml *Taq* DNA Polymerase supplied in a proprietary reaction buffer (pH 8.5), 400 μM of dATP, dGTP, dCTP, and dTTP, in addition to 3 mM of MgCl₂. All procedures were carried out in a sterile and stable environment to prevent external contamination.

PCR was undertaken in a thermal cycler (MJ, U.S.A.) with a mixture containing 20 units of nuclease-free water, 25 units of PCR Master Mix, 0.5 units of each designed cytokine genes primer (Invitrogen, USA) (Table 2 & 3). All primers were designed using the Roche UPL Primer Design Program, and 3 units of the extracted DNA sample were mixed to undergo thermal cycling. The thermal cycler was applied for PCR amplification of the DNA samples according to conditions set out in Table 2 [13,23,31]. All products from thermal cycling were labeled accordingly and stored at -80°C until use. In this study, β-actin was used (Table 2) with the forward primer 5'-CCTCTATGCCAACACAGTGC-3' and reverse primer 5'-ATACTCCTGCTTGCTGATCC-3'. β-actin is considered to be a constitutive housekeeping gene for PCR, having been used to compare changes in specific gene expressions [51].

Table 1 The clinical data (Mean ± SD) of control subjects and periodontitis patients

Parameters	Control subjects (N = 850)	Periodontitis patients (N = 440)
Age (years)	42.9 ± 9.7	49.3 ± 13.6
Age range (years)	26 - 60	28 - 65
Male / female	544 / 306	260 / 180
PD (mm)	2.7 ± 1.2	6.1 ± 2.7*
Sites% with BOP	40.3 ± 9.5	78.2 ± 19.8*
Sites% with gingival recession	1 ± 1.2	38.9 ± 25.9*
Sites% with calculus	34.1 ± 13.6	63.0 ± 25.8
CAL (mm)	0.0	6.2 ± 2.9

PD: probing depth; BOP: bleeding on probing; CAL: clinical attachment loss. Significant difference from the control subjects, **p* < 0.05.

Table 2 PCR conditions for various genes

Gene	Product size (bp)	Denaturation	Annealing	Extension	Cycles
IL-1 α	229	94°C, 1min	55°C, 30s	72°C, 60s	35
IL-1 β	305	94°C, 5mins	56°C, 45s	72°C, 60s	35
IL-6	296	95°C, 60s	60°C, 60s	72°C, 60s	35
TNF- α	133	94°C, 1min	61°C, 1min	72°C, 60s	35
IFN- γ	366	95°C, 5mins	56°C, 30s	72°C, 5mins	30
IL-4	195	95°C, 5mins	51°C, 60s	72°C, 60s	35
IL-10	139	94°C, 30s	60°C, 45s	72°C, 60s	35
β -actin	211	94°C, 30s	55°C, 30s	72°C, 60s	30

Restriction digest using *Fnu4H1*, *Aval*, *Nla III*, *Alw*, *Mnl I* and *AvaII* for IL-1 α , IL-1 β , IL-6, IFN- γ , IL-10, and TNF- α and IL-4 respectively

The amplified fragments generated from the PCR procedure were digested: 1) the 229 bp fragment on IL-1 α was recognized by *Fnu4H1* (Fermentas Life Sciences, U.S.A.) [52], 2) the 305 bp fragment on IL-1 β was recognized by *Aval* (Fermentas Life Sciences, U.S.A.) [53], 3) the 296 bp fragment on IL-6 was recognized by *NlaIII* (Fermentas Life Sciences, U.S.A.), [54] 4) the 296 bp fragment on IFN- γ was recognized by *Alw* (Fermentas Life Sciences, U.S.A.) [55], 5) the 139 bp fragment on IL-10 was recognized by *Mnl I* (Fermentas Life Sciences, U.S.A.) [56], and 6) both the 296 bp fragment on TNF- α and 195 bp fragment on IL-4 were recognized by *AvaII* (Fermentas Life Sciences, U.S.A.) [57,58].

For each digest, 10 μ l of the amplified PCR product was mixed with 2.5 to 5 units of the corresponding restriction enzyme, 10 μ l of nuclease-free water and 0.5

to 1 μ l of restriction enzyme buffer. The entire mixture was incubated for more than 4 hours at 37°C (Table 3) [13,23,31]. All digestion reagents were kept on ice before incubation to prevent denaturation. Every sample was digested twice to ensure for consistency of the amplicons.

Electrophoresis and visualization of digest product

The 10 μ l of digestion product and 1 μ l of Ready-Load 1 Kb DNA Ladder (Invitrogen, Spain) were loaded into 2-4% agarose gel (Invitrogen, Spain) containing 0.5 μ g/ml of ethidium bromide [13,23,31]. The gel underwent electrophoresis at 100 volts and 100 milliAmperes for 30 minutes. Afterward, the gel was visualized using a Dolphin-DOC ultraviolet illuminator (Wealtec, South Africa).

Sera measured by ELISA

ELISA was performed using the serum samples, following the manufacturer's instructions from cytokines ELISA Kit (Diacolone, France) [59]. The normal detection ranges of biomarkers are detailed in Table 4. The procedures were as follows: 100 μ l of the standard group solutions and serum of each subject were pipetted into a 96-well microplate. The plate was incubated for 2 to 3 hours at 350 rpm and washed with washing buffer three times. Then the wells were dried and 200 μ l of substrate tetramethylbenzidine was added into each well for 20 min in the dark at room temperature. The plates were read at 450 nm wavelength using Universal Microplate Reader (Sunrise, TECAN, Austria). The levels of cytokines in the samples were obtained by comparison with the standard curve generated from standards supplied by the manufacturer [59].

Table 3 The primer sequences and restriction enzymes used for detection of cytokine DNA polymorphism genes

Cytokines	Primers	Sequence	Position	Restriction Enzyme	Digestion Time (hours)	References
IL-1 α	Forward	5'ATGGTTTATAGAAATCATCAAGCCTAGGCA-3'	-889	<i>Fnu4H1</i>	>12	Walker et al.
	Reverse	5'AATGAAAGGAGGGGAGGATGACAGAAATGA-3'				
IL-1 β	Forward	5'-TGGCATTGATCTGGTTCATC-3'	-511	<i>Aval</i>	>12	Néstor et al.
	Reverse	5'-GTTTAGGAATCTTCCCACTT-3'				
IL-6	Forward	5'TTGTCAGACATGCCAAGTGCT-3'	-174	<i>Nla III</i>	4	Trevilatto et al.
	Reverse	5'-GCCTCAGAGACATCTCCAGTCC-3'				
TNF- α	Forward	5'-GAAGCCCTCCCAAGTCTAGT TC-3'	-238	<i>AvaII</i>	4	Sleijffers et al.
	Reverse	5'-CACTCCCCATCCTCCCTGGTC-3'				
IFN- γ	Forward	5'-GCTGTCATAATAATATTCAGAC-3'	-874	<i>Alw</i>	4	Inoue et al.
	Reverse	5'-CGAGCTTTAAAGATAGTTCC-3'				
IL-4	Forward	5'-TAAACTTGGGAGAACATGGT-3'	-590	<i>Ava II</i>	>12	Scarel-Caminaga et al.
	Reverse	5'-TGGGGAAAGATAGAGTAATA-3'				
IL-10	Forward	5'-CTCGTGCAACCCAACTGGC-3'	-1082	<i>Mnl I</i>	4	Chin et al.
	Reverse	5'-TCTTACCTATCCCTACTTCC-3'				

Table 4 Complete blood count (Mean \pm SD) of control subjects (N = 850) and periodontitis patients (N = 440)

Parameters	Control subjects	Periodontitis patients	Normal Range	Unit
White blood cell	6.33 (\pm 1.31)	3.96 (\pm 1.02)	4.00 - 11.00	10 ⁹ /L
Red blood cell	4.52 (\pm 0.13)	4.25 (\pm 0.10)	3.8 - 6.0	10 ¹² /L
Hemoglobin	15.0 (\pm 0.50)	12.5 (\pm 0.43)	11.5 - 16.5	g/dL
Platelet	273 (\pm 27.12)	241 (\pm 23.31)	150 - 400	10 ⁹ /L
Neutrophils	4.45 (\pm 1.05)	*1.68 (\pm 0.71)	2.0 - 7.5	10 ⁹ /L
Lymphocytes	1.37 (\pm 0.64)	*1.42 (\pm 0.26)	1.30 - 3.5	10 ⁹ /L
Monocyte	0.34 (\pm 0.19)	0.18 (\pm 0.11)	0.2 - 0.7	10 ⁹ /L
Eosinophil	0.12 (\pm 0.60)	0.06 (\pm 0.02)	0.0 - 0.5	10 ⁹ /L
Basophil	0.05 (\pm 0.03)	0.02 (\pm 0.01)	0.0 - 0.1	10 ⁹ /L
Neutrophils	70.3 (\pm 13.42)	*49.9 (\pm 11.43)	4.0 - 75.0	%
Lymphocytes	21.6 (\pm 9.06)	*42.3 (\pm 10.69)	20 - 45	%
Monocyte	5.4 (\pm 3.02)	5.4 (\pm 2.89)	4.7 - 12.2	%
Eosinophil	1.9 (\pm 0.34)	1.8 (\pm 0.52)	0.7 - 7.0	%
Basophil	0.8 (\pm 0.12)	0.6 (\pm 0.24)	0.1 - 1.2	%

Significant difference from the control, * $p < 0.05$

Statistical analysis

The Chi-squared test was applied to examine the differences in genotype distribution, allele frequency and carriage rate between healthy and patient groups. The alleles were calculated as an odds ratio (OR) with 95% confidence interval (95% CI) [13,23,31]. The soluble protein levels of cytokines were evaluated with an independent t-test [59], and $p < 0.05$ was considered statistically significant. Statistical analysis was performed using SPSS 18.0 for windows (SPSS Inc., U.S.A.).

Results

The clinical parameters of the healthy subjects and periodontitis patients were shown in Table 1. The periodontitis patient group exhibited a significantly greater mean of PD (6.1 ± 2.7 mm vs. 2.7 ± 1.2 mm), CAL (6.2 ± 2.9 mm vs 0.0) and a higher percentage of sites with BOP ($78.2 \pm 19.8\%$ vs. $40.3 \pm 9.5\%$) and gingival recession ($38.9 \pm 25.9\%$ vs. $1 \pm 1.2\%$) than the control group ($p < 0.05$). There was no significant difference found in the age and gender ratio between the two groups.

The total blood count of the healthy control subjects and periodontitis patients are presented in Table 4. With regard to the clinical parameters, the results of all

subjects were within the normal range, although the patient group exhibited a relatively higher count and percentage of lymphocytes, and a relatively lower count and percentage of neutrophils ($p < 0.05$) compared to the periodontally healthy controls.

The protein expression of patient and control samples for IL-1 α , IL-1 β , IL-6, TNF- α , IFN- γ , IL-10, and IL-4 (Table 5) measured by ELISA confirmed a statistically significant difference ($p < 0.001$). The digestion of fragments of various genes examined under a gel documentation system (Wealtec, South Africa) showed that the pro-inflammatory cytokines IL-1 α and TNF- α , and the anti-inflammatory cytokines IL-4 and IL-10 demonstrated an association with chronic inflammation in patients (X^2 : $p < 0.0001$).

After digestion by *Fnu4HI*, IL-1 α formed the DNA products of 153 and 76 bp for homozygous C allele, and 124, 76 and 29 bp for homozygous T alleles [52]. The detection frequency of the homozygous C allele of IL-1 α was similar in the subjects with chronic periodontitis (62%) and control group (55%) (X^2 , $p = 0.18$), but in the C/C genotype, the periodontitis patients presented a frequency of 52%, compared to 20% in the control group. The odds ratio for carriage of IL-1 α allele (T/T + C/T

Table 5 Protein expression (Mean \pm SD) of various cytokines in patient and control groups as measured by ELISA

Cytokines	Control subjects (N = 850)	Periodontitis patients (N = 440)	p value	Detection Range pg/ml
IL-1 α	24.08 (\pm 4.62)	59.73 (\pm 11.36)	0.0008	31.25 - 1000
IL-1 β	19.57 (\pm 3.75)	78.07 (\pm 17.09)	0.0004	15.6 - 500
IL-6	27.35 (\pm 2.87)	92.94 (\pm 17.44)	0.0005	2 - 200
TNF- α	26.90 (\pm 4.55)	67.61 (\pm 14.60)	0.0002	8 - 800
IFN- γ	52.77 (\pm 8.12)	94.71 (\pm 9.90)	0.0003	5 - 400
IL-4	1.16 (\pm 0.20)	4.62 (\pm 0.58)	0.0003	0.7 - 35
IL-10	22.70 (\pm 3.41)	46.89 (\pm 5.85)	0.0012	5 - 400

genotypes combined to compare with the C/C genotype) was 4.368 (95% CI = 2.309 - 8.264, X^2 : $p < 0.0001$) in periodontitis patients (Table 6).

After digestion by *AvaII*, TNF- α formed the DNA products of 70 and 63 bp for homozygous G alleles, and 63, 49 and 21 bp for homozygous A alleles.⁸¹ The detection frequency of the homozygous G allele of TNF- α was higher in the subjects with chronic periodontitis (57%) than the control group (34%) (X^2 : $p < 0.0001$, OR: 2.505, CI = 1.606 - 3.910). Comparing the G/G genotype, periodontitis patients presented higher a frequency of 43% than 17% in the healthy controls ($p < 0.0001$). The odds ratio for carriage of TNF- α allele (A/A and A/G genotypes combined to compare with the G/G genotype) was

3.716 (95% CI = 1.943 - 7.104) in periodontitis patients (Table 6).

The homozygous T alleles of IL-4 were represented by a DNA band with a size of 195 bp, homozygous C alleles were represented by DNA bands with sizes of 18 and 177 bp [58]. The detection frequency of the homozygous T alleles of IL-4 was higher in the patient (76%) than control group (45%). The Chi-Square test result was $p < 0.0001$, and the odds ratio was 3.837 (95% CI = 2.308 - 6.380) between the patient and control group. Comparing T/T genotype, the periodontitis patients presented a higher frequency of 55% in contrast to 18% in periodontally healthy controls ($p < 0.001$). The odds ratio for carriage of IL-4 allele (T/T and C/T genotypes combined compared with

Table 6 Genotype and allele frequency of cytokines in periodontitis patients (CP) and control subjects

Genotype and allele frequency of cytokines in periodontitis patients (CP) and control subjects									
Genotypes	CP Patients <i>n</i> = 440 (%)	Healthy subjects <i>n</i> = 850 (%)	CP versus Controls		Alleles	CP patients <i>n</i> = 880 (%)	Healthy subjects <i>n</i> = 1700 (%)	CP versus Controls	
			OR (95% CI)	χ^2 <i>p</i> values				OR (95% CI)	χ^2 <i>p</i> values
IL-1α									
*C/C	229(52)	170(20)	4.3681	23.32	C	546(62)	935(55)	1.13597	1.78
T/T	123(28)	85(10)	(2.3089-8.2638)	<0.0001	T	334(38)	765(45)	(0.865-2.1374)	0.1822
C/T	88(20)	595 (70)							
IL-1β									
*T/T	128(29)	314(37)	0.7132	0.97	T	510(58)	986(58)	0.9974	0
C/C	61(14)	179(21)	(0.3624-1.4035)	0.3247	C	370(42)	714(42)	(0.6393-1.5559)	1.000
C/T	251(57)	357(42)							
IL-6									
*G/G	110(25)	408(48)	0.3595	8.57	G	510(58)	1037(61)	0.8781	0.33
C/C/	25(8)	221(26)	0.177-0.7301	0.0034	C	370(42)	663(39)	(0.5626-1.3706)	0.5657
G/C	295(67)	221(26)							
TNF-α									
*G/G	189(43)	145(17)	3.7156	17.47	G	502(57)	578(34)	2.5054	17.23
A/A	132(30)	408(48)	(1.9432-7.1043)	0.0001	A	378(43)	1122(66)	(1.6056-3.9095)	0.0001
A/G	119(27)	297(35)							
IFN-γ									
*T/T	251(57)	476(56)	1.0382	0.010.9203	T	580(66)	1105(65)	1.0475	0.04
A/A	106(24)	221(26)	(0.5548-1.9427)		A	300(34)	595(35)	(0.6594-1.664)	0.8415
T/A	83(19)	153(18)							
IL-4									
*C/C	9(2)	230(27)	16.0032	13.2	C	211(24)	935(55)	3.8369	30.16
T/T	242(55)	153(18)	(2.1784-117.5633)	0.0003	T	669(76)	765(45)	(2.3075-6.3797)	0.0001
T/C	189(43)	467(55)							
IL-10									
*A/A	277(63)	782(92)	6.5546	5.211	A	590(67)	1581(93)	6.5731	63.26
G/G	132(30)	51(6)	(3.2316-13.2948)	0.0001	G	290(33)	119(7)	(3.9186-11.0259)	0.0001
A/G	31(7)	17(2)							

IL-1 α : T/T+C/T versus C/C; IL-1 β : C/C+ C/T versus T/T;

IL-6: C/C+G/C versus G/G; TNF- α : A/A+A/G versus G/G;

IFN- γ : A/A+T/A versus T/T; IL-4: T/T +T/C versus C/C;

IL-10: G/G+A/G versus A/A

OR, odds ratio; CI, confidence interval;

Significant difference from the control, * $p < 0.05$

the C/C genotype) was 16.003 (95% CI = 2.178 - 117.563) in periodontitis patients (Table 6).

The homozygous A alleles of IL-10 were represented by a DNA band with a size of 139 bp, and homozygous G alleles were represented by DNA bands with sizes of 106 and 33 bp. The detection frequency of the homozygous A alleles of IL-10 was higher in periodontally healthy controls (93%) than the patient group (67%). The result of the Chi-square test was $p < 0.001$, and the odds ratio was 6.573 (95% CI = **3.919** - 11.026) between the disease and control group. Comparing A/A genotype, the periodontitis patients presented lower frequency of 63% than 92% in control ($p < 0.001$). The odds ratio for carriage of IL-10 allele (G/G and A/G genotypes combined compared with the A/A genotype) was 6.555 (95% CI: 3.232-13.265) in periodontitis patients (Table 6).

Other genes investigated in the patients and healthy subjects (IL-1 β , IL-6, IFN- γ and IL-10) did not show any significant difference ($p > 0.05$), and the lower bound of confidence interval was below 1.

Overall, significant difference was found in the distribution of IL-1 α , TNF- α , IL-4 and IL-10 between the two groups (Table 6). For IL-1 α , the detection frequency of C allele was 62% in the patients group and 55% in the control. The subjects with the C allele are at similar risk for both groups. The genotype of C/C was significantly higher in the patient group (52%) than in the controls (20%) ($p < 0.001$). Individuals with the C/C genotype are at over four times higher risk for moderate to severe periodontitis than people with T/T and C/T genotypes ($p < 0.001$, OR = 4.368 with 95% CI 2.309 - 8.264). The detection frequency of G allele in TNF- α was 57% in the patients group and 34% in the control group. The patients with the G allele are at two-fold risk for the disease. The genotype of G/G was significantly higher in the patient group (43%) than in the controls (17%) ($p < 0.001$). Individuals with the G/G genotype are at nearly four times higher risk for moderate to severe periodontitis than persons with A/A and A/G genotypes ($p < 0.001$, OR = 3.716 with 95% CI 1.943 - 7.104). In IL-4, the detection frequency of C allele was 24% in the patient group and 55% in the healthy control group. The control subjects with the C allele are at almost four times less risk for the disease. The genotype of C/C was significantly lower in the patients (2%) compared to the healthy controls (27%) ($p < 0.001$). Individuals with the C/C genotype are at over sixteen times higher risk for moderate to severe periodontitis than persons with T/T and C/T genotypes ($p < 0.001$, OR = 16.003 with 95% CI = 2.178 - 117.563). The detection frequency of A allele in IL-10 was 67% in the patient group and 93% in the healthy control group. The subjects with the A allele are at six-fold less risk to the disease. The genotype of A/A was significantly lower in the patients group (63%) than in the controls (92%) ($p < 0.001$). Individuals with the A/A

genotype are over six times at higher risk for moderate to severe periodontitis than persons with G/G and A/G genotypes ($p < 0.001$, OR = 6.555 with 95% CI = 3.232 - 13.295).

Discussion

The levels of the proinflammatory cytokines IL-1 α , IL-1 β , TNF- α , IFN- γ and IL-6 are characteristically increased in diseased periodontal tissues or gingival crevicular fluid [60-63]. On the contrary, the initiation and progression of periodontal inflammation may be due to a lack or inappropriate response of the anti-inflammatory cytokines IL-4 and IL-10 in chronic periodontitis [61-63]. The role of polymorphisms in host responses in the progression of periodontitis has been well documented in scientific literature. In most situations, the genetic polymorphisms cause a change in the protein or its expression altering the immune response.

The results presented here demonstrate a similar frequency of C and T alleles of IL-1 α -889 in CP and control groups. The frequency of the C allele in CP patients was 62% which is comparable to other studies in some populations [64]. The latter demonstrates an important issue that genetic polymorphism vary among different ethnic populations, because the carriage rate of C allele tend to be lower in other Asian populations than the Chinese subjects in our study [65,66]. The frequency of T allele, in contrast, was higher in the control than CP group. The IL-1 α CC genotype (OR = 4.368; 95% CI = 2.309 to 8.264) was significantly associated with CP. This is in accordance with other similar studies [67]. In this study, IL-1 β -511 failed to demonstrate an association with periodontitis, which corresponds to four other studies to date [68,69]. The carriage rate of rare allele was the same in the patient and control groups. In all other previous reports, this rate was similar.

Another proinflammatory cytokine, TNF- α , possesses a wide range of immunoregulatory functions, including production of secondary mediators [70]. Many studies have investigated the SNPs in the promoter region at positions -1031, -863, -857, -376, -308, and -238. Among the above polymorphic sites, position -308 has been most researched, although most studies failed to demonstrate an association with periodontitis. However, the present study managed to showed an association of TNF- α GG genotype (OR = 3.716; 95% CI = 1.943 - 7.104) with periodontitis. Our studied cases showed a higher frequency of homozygous (G/G) genotype compared to healthy controls, which could be considered a risk genotype for periodontitis susceptibility.

In the current study, IL-6 did not exhibit an association with periodontitis, in contrast to studies on Caucasians and Brazilians [31,37]. The carriage rate of both C and G alleles was similar in CP and control groups. However,

the total cases showed a significantly higher frequency of heterozygous genotype IL-6 -174 (G/C) compared to the periodontally healthy controls. Thus, IL-6 -174 G/C genotype may be considered a risk genotype for periodontitis susceptibility.

The IL-4 -590 promoter polymorphism is the most studied polymorphisms of IL-4. Despite so, case-control studies have not shown any relationship between the IL-4 gene polymorphism and susceptibility to chronic periodontitis [71,72]. In contrast, our results demonstrated an association of IL-4 C/C genotype (OR = 16.003; 95% CI = 2.178 - 117.563) with a lower frequency in periodontitis patients. The T/T genotype, however, was more prevalent in our patients with periodontitis (55%), suggesting that this may be a risk genotype for periodontitis susceptibility. A reduction in the frequency of the C/C genotype in patients confers a protective influence against the development of the disease.

The IL-10 -1082 polymorphism was evident in our results, displaying a lower frequency of the A/A genotype in periodontitis patients than the healthy controls (63% versus 92%) [32-34]. A higher frequency of IL-10 -1082 G/G genotype was found in periodontitis patients (30%), compared to the periodontally healthy controls (6%). The results of the clinical trial were in accordance with previous reports: the proportion of subjects exhibiting the G/G genotype was significantly higher in subjects with severe periodontitis than in periodontally healthy individuals [73]. The incongruous results can be explained by the racial difference or association with other interactive types of cytokines or genetic markers predisposing for the disease in the studied cases.

Conclusions

The cytokine gene polymorphisms may be used as a marker for periodontitis susceptibility, clinical behaviour and severity. This detection offers early diagnosis and induction of prophylaxis to other family members against disease progression.

Acknowledgements

This article has been published as part of *Journal of Translational Medicine* Volume 10 Supplement 1, 2012: Selected articles from the Organisation for Oncology and Translational Research (OOTR) 7th Annual Conference. The full contents of the supplement are available online at <http://www.translational-medicine.com/supplements/10/S1>.

Author details

¹UNIMED Medical Institute, Hong Kong SAR. ²School of Chinese Medicine, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong SAR. ³Stomatological Hospital of Guangzhou Medical College, Guangzhou, PRC. ⁴Department of Stomatology, Sichuan Academy of Medical Sciences & Sichuan Provincial People's Hospital, No.32, Section 2, 1st Ring Road (West), Chengdu, Sichuan, PRC. ⁵State Key Laboratory for Oral Diseases and Department of Prosthodontics, West China Hospital of Stomatology, Sichuan University, Sichuan, PRC. ⁶Jin Hua Dentistry, Chengdu, 610041, Sichuan, PRC.

Authors' contributions

WTYL conducted the research, performed data collection and data analysis, and participated in manuscript writing. YY and CF, JL and YT conducted the clinical examination and performed data collection. LB performed data collection and data analysis. MW supervised clinical examination and participated in manuscript planning. HL performed data analysis and participated in manuscript writing. MNBC conducted the clinical examination and participated in manuscript writing. LWCC participated in manuscript planning and writing. Dr. Liu Qing was responsible for data collection. All authors read and approved the final manuscript.

Competing interests

We declare that we have no financial and personal relationships with other people or organizations that can inappropriately influence our work, there is no professional or other personal interest of any nature or kind in any product, service and/or company that could be construed as influencing the position presented in, the article entitled, "Gene polymorphism and protein of human pro- and anti-inflammatory cytokines in Chinese healthy subjects and chronic periodontitis patients".

Published: 19 September 2012

References

- Genco RJ: Host responses in periodontal diseases: current concepts. *J Periodontol* 1992, **63**:338-355.
- American Academy of Periodontology: Parameters of Care Supplement: Parameter on Periodontitis Associated with Systemic Conditions. *J Periodontol* 2000, **71**(5 Suppl):847-83.
- Michalowicz B: Genetic and heritable risk factors in periodontal disease. *J Periodontol* 1994, **65**:479-488.
- Hart TC, Kornman KS: Genetic factors in the pathogenesis of periodontitis. *Periodontology* 2000 1997, **14**:202-215.
- Elenkov IJ, Iezzoni DG, Daly A, Harris AG, Chrousos GP: Cytokine dysregulation, inflammation and well-being. *Neuroimmunomodulation* 2005, **12**:255-69.
- Malkowska A, Kasprzak A, Stopa J: Proinflammatory cytokines in pathogenesis of periodontal disease. *Pol Merkur Lekarski* 2006, **20**:93-8.
- Prasanna SJ, Gopalakrishnan D, Shankar SR, Vasandan AB: Pro-inflammatory cytokines, IFN γ and TNF α , influence immune properties of human bone marrow and Wharton jelly mesenchymal stem cells differentially. *PLoS One* 2010, **5**:e9016.
- Yano S, Sone S, Nishioka Y, Mukaida N, Matsushima K, Ogura T: Differential effects of anti-inflammatory cytokines (IL-4, IL-10 and IL-13) on tumoricidal and chemotactic properties of human monocytes induced by monocyte chemotactic and activating factor. *J Leukoc Biol* 1995, **57**:303-9.
- Gabay C, Lamacchia C, Palmer G: IL-1 pathways in inflammation and human diseases. *Nat Rev Rheumatol* 2010, **6**:232-41.
- McDevitt MJ, Wang HY, Knobelman C, Newman MG, di Giovine FS, Timms J, Duff GW, Kornman KS: Interleukin-1 genetic association with periodontitis in clinical practice. *J Periodontol* 2000, **71**:156-163.
- Cullinan MP, Westerman B, Hamlet SM, Palmer JE, Faddy MJ, Lang NP, Seymour GJ: A longitudinal study of interleukin-1 gene polymorphisms and periodontal disease in a general adult population. *J Clin Periodontol* 2001, **28**:1137-1144.
- Shirodaria S, Smith J, McKay IJ, Kennet CN, Hughes FJ: Polymorphisms in the IL-1A gene are correlated with levels of interleukin-1 α protein in gingival crevicular fluid of teeth with severe periodontal disease. *J Dent Res* 2000, **79**:1864-1869.
- Nikolopoulos GK, Dimou NL, Hamodrakas SJ, Bagos PG: Cytokine gene polymorphisms in periodontal disease: a meta-analysis of 53 studies including 4178 cases and 4590 controls. *J Clin Periodontol* 2008, **35**:754-67.
- Shete AR, Joseph R, Vijayan NN, Srinivas L, Banerjee M: Association of single nucleotide gene polymorphism at interleukin-1 β +3954, -511, and -31 in chronic periodontitis and aggressive periodontitis in Dravidian ethnicity. *J Periodontol* 2010, **81**:62-9.
- Liz-Graña M, Gómez-Reino Carnota J: Tumour necrosis factor. Genetics, cell action mechanism and involvement in inflammation. *Allergol Immunol Clin* 2001, **16**:140-149.

16. Brinkman BM, Kaijzel EL, Huizinga TW, Giphart MJ, Breedveld FC, Verweij CL: **Detection of a C-insertion polymorphism within the human tumor necrosis factor alpha (TNFA) gene.** *Hum Genet* 1995, **96**:493.
17. D'Alfonso S, Richiardi PM: **A polymorphic variation in a putative regulation box of the TNFA promoter region.** *Immunogenetics* 1994, **39**:150-154.
18. Ugliarolo AM, Turbay D, Pesavento PA, Delgado JC, McKenzie FE, Gribben JG, Hartl D, Yunis EJ, Goldfeld AE: **Identification of three new single nucleotide polymorphisms in the human tumor necrosis factor-alpha gene promoter.** *Tissue Antigens* 1998, **52**:359-367.
19. Wilson AG, di Giovine FS, Blakemore AI, Duff GW: **Single base polymorphism in the human tumour necrosis factor alpha (TNF alpha) gene detectable by NcoI restriction of PCR product.** *Hum Mol Genet* 1992, **1**:353.
20. Louis E, Franchimont D, Piron A, Gevaert Y, Schaaf-Lafontaine N, Roland S, Mahieu P, Malaise M, De Groote D, Louis R, Belaiche J: **Tumour necrosis factor (TNF) gene polymorphism influences TNF-alpha production in lipopolysaccharide (LPS)-stimulated whole blood cell culture in healthy humans.** *Clin Exp Immunol* 1998, **113**:401-406.
21. Fassmann A, Holla LI, Buckova D, Vasku A, Znojil V, Vanek J: **Polymorphisms in the +252(A/G) lymphotoxin-alpha and the -308(A/G) tumor necrosis factor-alpha genes and susceptibility to chronic periodontitis in a Czech population.** *J Periodontol Res* 2003, **38**:394-399.
22. Kornman KS, di Giovine FS: **Genetic variations in cytokine expression: a risk factor for severity of adult periodontitis.** *Ann Periodontol* 1998, **3**:327-38.
23. Endo M, Tai H, Tabeta K, Kobayashi T, Yamazaki K, Yoshie H: **Analysis of single nucleotide polymorphisms in the 5'-flanking region of tumor necrosis factor-alpha gene in Japanese patients with early-onset periodontitis.** *J Periodontol* 2001, **72**:1554-9.
24. Perez C, Gonzalez FE, Pavez V, Araya AV, Aguirre A, Cruzat A, Contreras-Levicoy J, Dotte A, Aravena O, Salazar L, Catalan D, Cuenca J, Ferreira A, Schiattino I, Aguillon JC: **The -308 polymorphism in the promoter region of the tumor necrosis factor-alpha (TNF-alpha) gene and ex vivo lipopolysaccharide-induced TNF-alpha expression in patients with aggressive periodontitis and/or type 1 diabetes mellitus.** *Eur Cytokine Netw* 2004, **15**:364-370.
25. Trevisatto PC, Tramontina VA, Machado MA, Goncalves RB, Sallum AW, Line SR: **Clinical, genetic and microbiological findings in a Brazilian family with aggressive periodontitis.** *J Clin Periodontol* 2002, **29**:233-239.
26. Craandijk J, van Krugten MV, Verweij CL, van der Velden U, Loos BG: **Tumor necrosis factor-alpha gene polymorphisms in relation to periodontitis.** *J Clin Periodontol* 2002, **29**:28-34.
27. Soga Y, Nishimura F, Ohshima H, Maeda H, Takashiba S, Murayama Y: **Tumor necrosis factor-alpha gene (TNF-alpha) -1031/-863, -857 single-nucleotide polymorphisms (SNPs) are associated with severe adult periodontitis in Japanese.** *J Clin Periodontol* 2003, **30**:524-31.
28. Holla LI, Fassmann A, Stejskalova A, Znojil V, Vanek J, Vacha J: **Analysis of the interleukin-6 gene promoter polymorphisms in Czech patients with chronic periodontitis.** *J Periodontol* 2004, **75**:30-36.
29. Nibali L, D'Aiuto F, Donos N, Griffiths GS, Parkar M, Tonetti MS, Humphries SE, Brett PM: **Association between periodontitis and common variants in the promoter of the interleukin-6 gene.** *Cytokine* 2009, **45**:50-4.
30. Komatsu Y, Tai H, Galicia JC, Shimada Y, Endo M, Akazawa K, Yamazaki K, Yoshie H: **Interleukin-6 (IL-6)-373 A9T11 allele is associated with reduced susceptibility to chronic periodontitis in Japanese subjects and decreased serum IL-6 level.** *Tissue Antigens* 2005, **65**:110-4.
31. Trevisatto PC, Scarel-Caminaga RM, de Brito RB Jr, de Souza AP, Line SR: **polymorphism at position -174 of IL-6 gene is associated with susceptibility to chronic periodontitis in a Caucasian Brazilian population.** *J Clin Periodontol* 2003, **30**:438-42.
32. D'Aiuto F, Parkar M, Brett PM, Ready D, Tonetti MS: **Gene polymorphisms in pro-inflammatory cytokines are associated with systemic inflammation in patients with severe periodontal infections.** *Cytokine* 2004, **28**:29-34.
33. D'Aiuto F, Ready D, Parkar M, Tonetti MS: **Relative contribution of patient-, tooth-, and site-associated variability on the clinical outcomes of subgingival debridement. I. Probing depths.** *J Periodontol* 2005, **76**:398-405.
34. Fishman D, Faulds G, Jeffery R, Mohamed-Ali V, Yudkin JS, Humphries S, Woo P: **The effect of novel polymorphisms in the interleukin-6 (IL-6) gene on IL-6 transcription and plasma IL-6 levels, and an association with systemic-onset juvenile chronic arthritis.** *J Clin Invest* 1998, **102**:1369-76.
35. Cesar-Neto JB, Duarte PM, de Oliveira MC, Casati MZ, Tambeli CH, Parada CA, Sallum EA, Nociti FH Jr: **Smoking modulates interferon-gamma expression in the gingival tissue of patients with chronic periodontitis.** *Eur J Oral Sci* 2006, **114**:403-408.
36. Pravica V, Perrey C, Stevens A, Lee JH, Hutchinson IV: **A single nucleotide polymorphism in the first intron of the human IFN-gamma gene: Absolute correlation with a polymorphic CA microsatellite marker of high IFNgamma production.** *Hum Immunol* 2000, **61**:863-866.
37. Babel N, Cherepnev G, Babel D, Tropmann A, Hammer M, Volk HD: **Analysis of tumor necrosis factor-alpha, transforming growth factor-beta, interleukin-10, IL-6, and interferon-gamma gene polymorphisms in patients with chronic periodontitis.** *J Periodontol* 2006, **77**:1978-83.
38. Sutherland GR, Baker E, Callen DF, Hyland VJ, Wong G, Clark S: **Interleukin 4 is at 5q31 and interleukin 6 is at 7p15.** *Hum Genet* 1988, **79**:335-337.
39. McFarlane CG, Meikle MC: **Interleukin-2, interleukin-2 receptor and interleukin-4 levels are elevated in the sera of patients with periodontal disease.** *J Periodontol Res* 1991, **26**:402-8.
40. Mout R, Willemze R, Landegent JE: **Repeat polymorphisms in the interleukin-4 gene (IL4).** *Nucleic Acids Res* 1991, **19**:3763.
41. Gonzales JR, Kobayashi T, Michel J, Mann M, Yoshie H, Meyle J: **Interleukin-4 gene polymorphisms in Japanese and Caucasian patients with aggressive periodontitis.** *J Clin Periodontol* 2004, **31**:384-389.
42. Pontes CC, Gonzales JR, Novaes AB Jr, Junior MT, Grisi MF, Michel J, Meyle J, de Souza SL: **Interleukin-4 gene polymorphism and its relation to periodontal disease in a Brazilian population of African heritage.** *J Dent* 2004, **32**:241-246.
43. Kim JM, Brannan CI, Copeland NG, Jenkins NA, Khan TA, Moore KW: **Structure of the mouse IL-10 gene and chromosomal localization of the mouse and human genes.** *J Immunol* 1992, **148**:3618-3623.
44. Scarel-Caminaga RM, Trevisatto PC, Souza AP, Brito RB, Camargo LE, Line SR: **Interleukin 10 gene promoter polymorphisms are associated with chronic periodontitis.** *J Clin Periodontol* 2004, **31**:443-448.
45. Eskdale J, Kube D, Tesch H, Gallagher G: **Mapping of the human IL10 gene and further characterization of the 5' flanking sequence.** *Immunogenetics* 1997, **46**:120-128.
46. Berglundh T, Donati M, Hahn-Zoric M, Hanson LA, Padyukov L: **Association of the -1087 IL 10 gene polymorphism with severe chronic periodontitis in Swedish Caucasians.** *J Clin Periodontol* 2003, **30**:249-54.
47. Machin D, Campbell MJ, Tan SB, Tan SH: **Sample Size Tables for Clinical Studies.** Wiley-Blackwell, 3 2009.
48. Lemeshow S, Hosmer DW, Klar J, Lwanga SK: **Adequacy of sample size in health studies Published on behalf of the World Health Organization.** Wiley-Blackwell 1990, 1-293.
49. Armitage GC: **Development of a classification system for periodontal diseases and conditions.** *Ann Periodontol* 1999, **4**:1-6.
50. Armitage GC: **Periodontal diagnoses and classification of periodontal diseases.** *Periodontology* 2000 2004, **34**:9-21.
51. Mori R, Wang Q, Danenberg KD, Pinski JK, Danenberg PV: **Both beta-actin and GAPDH are useful reference genes for normalization of quantitative RT-PCR in human FFPE tissue samples of prostate cancer.** *Prostate* 2008, **68**:1555-60.
52. Walker SJ, Van Dyke TE, Rich S, Kornman KS, di Giovine FS, Hart TC: **Genetic polymorphisms of the IL-1alpha and IL-1beta genes in African-American LJP patients and an African-American control population.** *J Periodontol* 2000, **71**:723-8.
53. Cox A, Camp NJ, Nicklin MJ, di Giovine FS, Duff GW: **An analysis of linkage disequilibrium in the interleukin-1 gene cluster, using a novel grouping method for multiallelic markers.** *Am J Hum Genet* 1998, **62**:1180-8.
54. Depboylu C, Lohmüller F, Gocke P, Du Y, Zimmer R, Gasser T, et al: **An interleukin-6 promoter variant is not associated with an increased risk for Alzheimer's disease.** *Dement Geriatr Cogn Disord* 2004, **17**:170-3.
55. Wang TN, Chu YT, Chen WY, Feng WW, Shih NH, Hsiang CH, et al: **Association of interferon-gamma and interferon regulatory factor 1 polymorphisms with asthma in a family-based association study in Taiwan.** *Clin Exp Allergy* 2006, **36**:1147-52.
56. Nemec P, Pavkova-Goldbergova M, Gatterova J, Fojtik Z, Vasku A, Soucek M: **Association of the -1082 G/A promoter polymorphism of interleukin-10 gene with the autoantibodies production in patients with rheumatoid arthritis.** *Clin Rheumatol* 2009, **28**:899-905.

57. Vinasco J, Beraún Y, Nieto A, Fraile A, Mataran L, Pareja E, et al: **Polymorphism at the TNF loci in rheumatoid arthritis.** *Tissue Antigens* 1997, **49**:74-8.
58. Scarel-Caminaga RM, Trevisatto PC, Souza AP, Brito RB Jr., Line SRP: **Investigation of IL-4 gene polymorphism in individuals with different levels of chronic periodontitis in a Brazilian population.** *J Clin Periodontol* 2003, **30**:341-345.
59. Loo WT, Jin LJ, Chow LW, Cheung MN, Wang M: **Rhodiola algida improves chemotherapy-induced oral mucositis in breast cancer patients.** *Expert Opin Investig Drugs* 2010, **19**(Suppl 1):S91-100.
60. Graves DT, Cochran D: **The contribution of interleukin-1 and tumor necrosis factor to periodontal tissue destruction.** *J Periodontol* 2003, **74**:391-401.
61. Stashenko P, Fujiyoshi P, Obernesser MS, Probstak L, Haffajee AD, Socransky SS: **Levels of interleukin 1 beta in tissue from sites of active periodontal disease.** *J Clin Periodontol* 1991, **18**:548-54.
62. Costa PP, Trevisan GL, Macedo GO, Palioto DB, Souza SL, Grisi MF, Novaes AB Jr, Taba M Jr: **Salivary interleukin-6, matrix metalloproteinase-8, and osteoprotegerin in patients with periodontitis and diabetes.** *J Periodontol* 2010, **81**:384-91.
63. Alexander MB, Damoulis PD: **The role of cytokines in the pathogenesis of periodontal disease.** *Curr Opin Periodontol* 1994, 39-53.
64. Laine ML, Farré MA, González G, van Dijk LJ, Ham AJ, Winkel EG, et al: **Polymorphisms of the interleukin-1 gene family, oral microbial pathogens, and smoking in adult periodontitis.** *J Dent Res* 2001, **80**:1695-9.
65. Anusakathien O, Sukboon A, Sitthiphong P, Teanpaisan R: **Distribution of interleukin-1beta(+3954) and IL-1alpha(-889) genetic variations in a Thai population group.** *J Periodontol* 2003, **74**:1796-802.
66. Kobayashi T, Ito S, Kuroda T, Yamamoto K, Sugita N, Narita I, et al: **The interleukin-1 and Fc-gamma receptor gene polymorphisms in Japanese patients with rheumatoid arthritis and periodontitis.** *J Periodontol* 2007, **78**:2311-8.
67. Wagner J, Kaminski WE, Aslanidis C, Moder D, Hiller KA, Christgau M, Schmitz G, Schmalz G: **Prevalence of OPG and IL-1 gene polymorphisms in chronic periodontitis.** *J Clin Periodontol* 2007, **34**:823-7.
68. Gore EA, Sanders JJ, Pandey JP, Palesch Y, Galbraith GM: **Interleukin-1beta +3953 allele 2: association with disease status in adult periodontitis.** *J Clin Periodontol* 1998, **25**:781-5.
69. Geismar K, Enevold C, Sørensen LK, Gyntelberg F, Bendtzen K, Sigurd B, et al: **Involvement of interleukin-1 genotypes in the association of coronary heart disease with periodontitis.** *J Periodontol* 2008, **79**:2322-30.
70. Sheng WS, Hu S, Ni HT, Rowen TN, Lokensgard JR, Peterson PK: **TNF-alpha-induced chemokine production and apoptosis in human neural precursor cells.** *J Leukoc Biol* 2005, **78**:1233-41.
71. Hooshmand B, Hajilooi M, Rafiei A, Mani-Kashani KH, Ghasemi R: **Interleukin-4 (C-590T) and interferon-gamma (G5644A) gene polymorphisms in patients with periodontitis.** *J Periodontol Res* 2008, **43**:111-5.
72. Holla LI, Fassmann A, Augustin P, Halabala T, Znojil V, Vanek J: **The association of interleukin-4 haplotypes with chronic periodontitis in a Czech population.** *J Periodontol* 2008, **79**:1927-33.
73. Berglundh T, Donati M, Hahn-Zoric M, Hanson LA, Padyukov L: **Association of the -1087 IL 10 gene polymorphism with severe chronic periodontitis in Swedish Caucasians.** *J Clin Periodontol* 2003, **30**:249-54.

doi:10.1186/1479-5876-10-S1-S8

Cite this article as: Loo et al.: Gene polymorphism and protein of human pro- and anti-inflammatory cytokines in Chinese healthy subjects and chronic periodontitis patients. *Journal of Translational Medicine* 2012 **10**(Suppl 1):S8.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

